AVIAX II - semduramicin powder

Phibro Animal Health

ACTIVE DRUG INGREDIENT(S)

(As semduramicin sodium biomass)

INGREDIENTS:

Dried semduramicin fermentation products, soybean mill run, mineral oil, sodium carbonate and sodium aluminosilicate.

FOR USE IN THE MANUFACTURE OF BROILER CHICKEN FEED.

CAUTION:

Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: avoid contact with skin, eyes, or mucous membranes; dust masks or respirators, safety glasses, gloves, and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions-seek prompt medical attention if such reactions are suspected.

IMPORTANT:

Must be thoroughly mixed with feed before use.

STORE AT OR BELOW 25°C (77°F)

SEE BACK OF PACKAGE FOR FURTHER USE DIRECTIONS

INDICATIONS FOR USE

For the prevention of coccidiosis in broiler chickens caused by Eimeria tenella, E. acervulina, E. maxima, E. brunetti, E. necatrix, and E. mitis.

MIXING AND FEEDING DIRECTIONS

Thoroughly mix 1 lb of Aviax II Type A Medicated Article per 2000 lb of feed ingredients to provide 25 ppm (22.7 g) of semduramicin per ton of Type C finished broiler feeds. Feed continuously as the sole ration to broiler chickens.

WARNING:

Do not feed to laying hens.

Aviax® is a registered trademark of Phibro Animal Health, Inc., Phibro Animal Health, Inc., Ridgefield Park, NJ 07660

Net Weight: 40 lb (18.2 kg)

NADA # 141-281, Approved by FDA 7950000 101-9091-03

TAKE TIME Made in USA 7950000







Net Weight: 40 lb (18.2 kg)

Phibre ANIMAL HEALTH







TYPE A MEDICATED ARTICLE

Indications for Use

For the prevention of coccidiosis in broiler chickens caused by Eimeria tenella, E. acervulina, E. maxima, E. brunetti, E. necatrix, and E. mitis.

Mixing and Feeding Directions
Thoroughly mix 1 lb of Aviax II Type A Medicated Article
per 2000 lb of feed ingredients to provide 25 ppm (22.7) gill of Semonduranich per ton of Type C finished broiler feeds. Feed continuously as the sole ration to broiler chickens.





Made in USA 7950000 101-9091-03



